

# Prevantics<sup>®</sup> range of applicators, swabsticks and swabs provide proven skin disinfection.


- + **Better than Traditional Methods:** the combination of 2% Chlorhexidine gluconate (CHG) and 70% Isopropyl alcohol (IPA) provides long-lasting microbial kill of up to 48 hours in comparison to 2 hours with just IPA alone.<sup>1</sup>
- + **Complies with Guidelines:** its 2% CHG and 70% IPA formulation is recommended in evidence-based infection guidelines published by health organisations such as NICE<sup>2</sup>, epic3<sup>3</sup> and the World Health Organization<sup>4</sup>.
- Proven Efficacy: based on its active ingredients, a broad spectrum of efficacy is guaranteed against bacterial, yeasticial & virucidal activity.







The single-use applicator is a ready to use device containing 2% (w/w) Chlorhexidine gluconate and 70% (v/v) Isopropyl alcohol requiring no activation.



## Recommended applicator based on general skin coverage

| Applicator Size | RECOMMENDED SKIN COVERAGE |
|-----------------|---------------------------|
| 3ml             | 15cm x 15cm               |
| 10.5ml          | 30cm x 30cm               |
| 26ml            | 50cm x 50cm               |

# Range Details:

|                  | REORDER NO | PRODUCT SIZE | QTY PER CASE   | CLASSIFICATION |
|------------------|------------|--------------|--|----------------|
| 26ml             | IAP4140    | 26ml         | 10 units per carton<br>6 cartons per case<br>60 units per case | Biocide        |
| 26ml with tint   | IAP4230    | 26ml         | 10 units per carton<br>6 cartons per case<br>60 units per case | Biocide        |
| 10.5ml           | IAP4130    | 10.5ml       | 10 units per carton<br>6 cartons per case<br>60 units per case | Biocide        |
| 10.5ml with tint | IAP4220    | 10.5ml       | 10 units per carton<br>6 cartons per case<br>60 units per case | Biocide        |
| 3ml              | IAP4110    | 3ml          | 11 units per carton<br>6 cartons per case<br>66 units per case | Biocide        |
| 3ml with tint    | IAP4200    | 3ml          | 11 units per carton<br>6 cartons per case<br>66 units per case | Biocide        |



The single-use, ready to use swabstick contains 2% (w/w) Chlorhexidine gluconate and 70% (v/v) Isopropyl alcohol requiring no activation.

Easy to open with no sharp glass to break.



The swabstick has been developed to allow the product to be safely removed from the packaging without any external contact, assisting aseptic non-touch technique (ANTT) compliance.

The swabstick has a unique 2-sided applicator for interdigital areas and skin folds ensuring thorough skin disinfection.

|                 | REORDER NO | PRODUCT SIZE | QTY PER CASE   | CLASSIFICATION |
|-----------------|------------|--------------|--|----------------|
| 1.6ml Swabstick | ISS4010    | 1.6ml        | 50 units per carton<br>10 cartons per case<br>500 units per case | Biocide        |







The ready to use, premoistened wipe contains 2% (w/w) Chlorhexidine gluconate (CHG) and 70% (v/v) Isopropyl alcohol (IPA) to help prevent the spread of infections and help improve patient outcomes.

|        | REORDER NO | PRODUCT SIZE  | QTY PER CASE  | CLASSIFICATION |
|--------|------------|---------------|---|----------------|
| Sachet | ISW4030    | 150mm x 129mm | 200 units per carton<br>8 cartons per case<br>1600 units per case | Biocide        |

# Other formulations available:











### References:

- <sup>1</sup> Analyses comparing the antimicrobial activity and safety of current antiseptic agents: A review, Hibbard JS, J Infusion Nursing 2005; 28: 194-207 & Skin antisepsis: past, present, and future, Crosby CT, Mares AK, J Vasc Access Devices 2001; Spring: 26-31 1.
- <sup>2</sup> National Institute for Clinical Excellence. Clinical Guideline 2. Infection Control.

Prevention of healthcare-associated infection in primary and community care. June 2003. http://www.nice.org.uk/pdf/ CG2fullguidelineinfectioncontrol.pdf

<sup>3</sup> epic3: National Evidence-Based Guidelines for Preventing Healthcare-Associated Infections in NHS Hospitals in England H.P. Lovedaya \*, J.A. Wilsona , R.J. Pratta , M. Golsorkhia , A. Tinglea , A. Baka , J. Brownea , J. Prietob, M. Wilcox. <sup>4</sup>World Health Organization. Global guidelines for the prevention of surgical site infection. Geneva, Switzerland. 2016. https://www. ncbi. nlm.nih.gov/books/NBK401132/pdf/Bookshelf\_ NBK401132.pdf

### Find out more about how our Layered Approach to Infection Prevention can help you.

PDI EMEA, Pywell Road, Willowbrook Est. Industrial Estate, Corby - Northants, NN17 5XJ, UK





+44 (0) 8081 697 945 Contact@pdi-emea.com wearepdi-intl.com in PDI International







